


My physician recommends...

Children's Advil®

Measure dosing only with the device that comes with the product



Based on weight and age, your healthcare professional has indicated below which Advil® medicine is right for your child.

Active Ingredients

Indications

Infants' Advil® Concentrated Drops
for ages 6-23 months

Ibuprofen 50 mg (NSAID)* in each 1.25 mL

Fever reducer, relieves minor aches and pains due to the common cold, flu, headaches, and toothaches

Children's Advil® Suspension
for ages 2-11 years

Ibuprofen 100 mg (NSAID)* in each 5 mL

Fever reducer, relieves minor aches and pains due to the common cold, flu, sore throat, headaches, and toothaches

Weight

Age

Patient Dosing

under 6 months

do not use

do not use

12-17 pounds

6-11 months

1.25 mL¹ every 6-8 hrs
not to exceed 4 doses/24 hrs

Use Infants' Advil® Concentrated Drops

18-23 pounds

12-23 months

1.875 mL¹ every 6-8 hrs
not to exceed 4 doses/24 hrs

1 tsp¹ every 6-8 hrs
not to exceed 4 doses/24 hrs

24-35 pounds

2-3 years

Use Children's Advil® Suspension

1½ tsp¹ every 6-8 hrs
not to exceed 4 doses/24 hrs

36-47 pounds

4-5 years

2 tsp¹ every 6-8 hrs
not to exceed 4 doses/24 hrs

48-59 pounds

6-8 years

2½ tsp¹ every 6-8 hrs
not to exceed 4 doses/24 hrs

60-71 pounds

9-10 years

3 tsp¹ every 6-8 hrs
not to exceed 4 doses/24 hrs

72-95 pounds

11 years

Use Advil® Tablets or Liqui-Gels®

Use Advil® Tablets or Liqui-Gels®

12+ years